



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

RHINOCILUS LECONTEI IN LOS
ANGELES.

In the COPEIA of February 20, 1915, (No. 15) appeared a list of fourteen snakes found in the vicinity of the city of Los Angeles. To those must now be added one more recently discovered at Belvidere, California, by Mr. Viggo Tarp. This is *Rhinochilus lecontei* (Baird & Girard) and was found on May 13, 1915, about two miles from the beach on a bare hill near some gopher holes. This specimen is now thriving in captivity on a diet of lizards. It is 38½ inches long.

The only other traces I have found of LeConte's Snake have been in La Canada Valley, where an old bee-man named Charlie Metcalf had skins of two other specimens found in the valley where he lives. Near Los Angeles, LeConte's Snake is considered rare.

PAUL D. R. RÜTHLING,
Los Angeles, California.